

On the Dynamical Generation of Quark-Level-Linear-Sigma-Model-like Theories beyond one Loop

Frieder Kleefeld ¹

*Collaborator of the
Centro de Física das Interações Fundamentais (CFIF), Instituto Superior Técnico,
Edifício Ciência, Piso 3, Av. Rovisco Pais, P-1049-001 LISBOA, Portugal*

Abstract. A self-consistent strategy to complete the dynamical generation of Quark-Level-Linear-Sigma-Model-like Lagrangean theories beyond one loop as proposed in more detail in our manuscript arXiv:0802.1540 [hep-ph] [1] is shortly outlined.

Keywords: dynamical generation, linear sigma model, quadratic divergencies, renormalization

PACS: 11.10.Cd, 11.10.Ef, 11.10.Gh, 12.40.-y

Throughout the construction of Lagrangean densities used e.g. in particle physics one faces at least two problems: 1) unproductive Lagrangeans contain too many uncorrelated parameters (masses, couplings) which have to be fitted to experiment; 2) inherent divergencies of logarithmic, linear, quadratic, ... type need to be renormalized. The concept of *dynamical generation* [1, 2] of Lagrangean theories addresses and solves both issues simultaneously:

- 1) In the spirit of Eguchi [3] one starts out from very few fundamental 3-point interaction vertices and constructs then on the basis of these vertices by “loop-shrinking” [2] the so-called effective action (and its underlying Lagrangean) containing also terms for all remaining n-point vertices between the fields making up the theory.
- 2) The couplings of the fundamental 3-point interaction vertices are then chosen such that linear, quadratic [4], ... divergencies cancel [1, 2] while the remaining logarithmic divergencies are renormalized [5] by adding to the effective action counter terms which replace [1] in the spirit of the log.-divergent gap equation [6, 7, 8] of Delbourgo and Scadron (DS) [6, 7] the integral $I_2(\overline{m}^2) \equiv \int \frac{d^4 p}{(2\pi)^4} \frac{1}{(p^2 - \overline{m}^2)^2}$ at some experimentally defined renormalization scale $\overline{m}$ (being in the case of DS approximately equal to the nonstrange constituent quark mass $\hat{m}$, i.e. $\overline{m} \simeq \hat{m} = m_q$) by some universal complex number $+ \frac{i}{16\pi^2}$.

For various reasons like e.g. the lacking [9, 10, 11] evidence for the existence of gluons and new developments in mathematical physics there has developed an alternative approach to strong interactions being different from QCD which is known since the

¹ e-mail: kleefeld@cfif.ist.utl.pt, URL: <http://cfif.ist.utl.pt/~kleefeld/>
Present address for postal correspondence: Pfisterstr. 31, 90762 Fürth, Germany

benchmarking work of DS [6, 7] as the Quark-Level Linear Sigma Model (QLL σ M).² Disregarding vector & axial vector mesons the SU(2) $\times$ SU(2) QLL σ M assuming $N_F = 2N_c = 6$ Fermions, one scalar isoscalar meson σ and $N_\pi = 3$ pions is constructed on the basis the interaction Lagrangean $\mathcal{L}_{\text{quark-meson}}(x) = g \bar{q}_+(x) (\sigma(x) + i \gamma_5 \vec{\tau} \cdot \vec{\pi}(x)) q_-(x)$ [6, 7, 14] yielding by loop-shrinking the leading terms in the Lagrangean of the effective action for meson-meson interactions, i.e. $\mathcal{L}_{\text{meson-meson}} = g_{\sigma\pi\pi} \sigma(x)(\sigma(x)^2 + \vec{\pi}(x)^2) - \frac{\lambda}{4} (\sigma(x)^2 + \vec{\pi}(x)^2)^2 + \dots$. Following the formalism described in Ref. [2] the relevant terms in the effective action of the SU(2) $\times$ SU(2) QLL σ M for the σ -one-point function (see Fig. 1), for the two-point function of the quarks (see Fig. 2), of the σ (see Fig. 3) and of the pions (see Fig. 4) are obtained³. After isolating the quadratically divergent part of the effective actions it is straight forward to extract the following two conditions which yield a complete cancellation of quadratic divergencies [1] :

² It has been the author to point out that the experimentally favoured asymptotically free phase of QLL σ M belongs to the acceptable class of non-Hermitian PT-symmetric field theories [2, 11, 12, 13].

³ The result is [1, 2] (NLTs = Non-Local Terms):

$$S_{(1)}[\sigma] = \int d^4x \sigma(x) i \left\{ -4g N_F m_q I_1(m_q^2) + 3g_{\sigma\pi\pi} I_1(m_\sigma^2) + N_\pi g_{\sigma\pi\pi} I_1(m_\pi^2) \right. \\ \left. + 2\lambda g_{\sigma\pi\pi} i \left(3 I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\sigma^2, m_\sigma^2) + N_\pi I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\pi^2, m_\pi^2) \right) \right\} + \text{NLTs}, \quad (1)$$

$$S_{(2)}[\bar{q}q] = \frac{i}{2} \int d^4x \bar{q}_+(x) q_-(x) \frac{2g}{m_\sigma^2} \left\{ -4g N_F m_q I_1(m_q^2) + 3g_{\sigma\pi\pi} I_1(m_\sigma^2) + N_\pi g_{\sigma\pi\pi} I_1(m_\pi^2) \right. \\ \left. + 2\lambda g_{\sigma\pi\pi} i \left(3 I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\sigma^2, m_\sigma^2) + N_\pi I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\pi^2, m_\pi^2) \right) \right\} \\ - \frac{i}{2} \int d^4x \bar{q}_+(x) q_-(x) 2g^2 m_q \left(I_{1,1}(m_q^2, m_\sigma^2) - N_\pi I_{1,1}(m_q^2, m_\pi^2) \right) + \text{NLTs}, \quad (2)$$

$$S_{(3)}[\sigma^2] = \frac{i}{2} \int d^4x \sigma(x)^2 \frac{6g_{\sigma\pi\pi}}{m_\sigma^2} \left\{ -4g N_F m_q I_1(m_q^2) + 3g_{\sigma\pi\pi} I_1(m_\sigma^2) + N_\pi g_{\sigma\pi\pi} I_1(m_\pi^2) \right. \\ \left. + 2\lambda g_{\sigma\pi\pi} i \left(3 I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\sigma^2, m_\sigma^2) + N_\pi I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\pi^2, m_\pi^2) \right) \right\} \\ + \frac{i}{2} \int d^4x \sigma(x)^2 4g^2 N_F \left(I_1(m_q^2) + 2m_q^2 I_2(m_q^2) \right) - \frac{i}{2} \int d^4x \sigma(x)^2 \lambda \left(3 I_1(m_\sigma^2) + N_\pi I_1(m_\pi^2) \right) \\ - \frac{i}{2} \int d^4x \sigma(x)^2 2\lambda^2 i \left(3 I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\sigma^2, m_\sigma^2) + N_\pi I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\pi^2, m_\pi^2) \right) \\ - \frac{i}{2} \int d^4x \sigma(x)^2 2g_{\sigma\pi\pi}^2 \left(9 I_2(m_\sigma^2) + N_\pi I_2(m_\pi^2) \right) + \text{NLTs}, \quad (3)$$

$$S_{(4)}[\vec{\pi}^2] = \frac{i}{2} \int d^4x \vec{\pi}(x)^2 \frac{2g_{\sigma\pi\pi}}{m_\sigma^2} \left\{ -4g N_F m_q I_1(m_q^2) + 3g_{\sigma\pi\pi} I_1(m_\sigma^2) \right. \\ \left. + N_\pi g_{\sigma\pi\pi} I_1(m_\pi^2) + 2\lambda g_{\sigma\pi\pi} i \left(3 I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\sigma^2, m_\sigma^2) + N_\pi I_{1,1,1}^{\text{sunset}}(m_\sigma^2, m_\pi^2, m_\pi^2) \right) \right\} \\ + \frac{i}{2} \int d^4x \vec{\pi}(x)^2 4g^2 N_F I_1(m_q^2) - \frac{i}{2} \int d^4x \vec{\pi}(x)^2 \lambda \left(I_1(m_\sigma^2) + (N_\pi + 2) I_1(m_\pi^2) \right) \\ - \frac{i}{2} \int d^4x \vec{\pi}(x)^2 2\lambda^2 i \left(I_{1,1,1}^{\text{sunset}}(m_\pi^2, m_\sigma^2, m_\sigma^2) + (N_\pi + 2) I_{1,1,1}^{\text{sunset}}(m_\pi^2, m_\pi^2, m_\pi^2) \right) \\ - \frac{i}{2} \int d^4x \vec{\pi}(x)^2 4g_{\sigma\pi\pi}^2 I_{1,1}(m_\sigma^2, m_\pi^2) + \text{NLTs}. \quad (4)$$

$0 = -4gN_F m_q + \left(1 - \frac{\lambda}{4\pi^2}\right) g_{\sigma\pi\pi}(3 + N_\pi)$ and $0 = +4g^2 N_F - \left(1 - \frac{\lambda}{4\pi^2}\right) \lambda (3 + N_\pi)$. These conditions can be finally solved for λ and $g_{\sigma\pi\pi}$ as a function of g and m_q [1]:

$$\lambda = 2\pi^2 \left(1 \pm \sqrt{1 - \frac{4g^2 N_F}{\pi^2(3 + N_\pi)}}\right), \quad g_{\sigma\pi\pi} = 2\pi^2 \left(1 \pm \sqrt{1 - \frac{4g^2 N_F}{\pi^2(3 + N_\pi)}}\right) \frac{m_q}{g}. \quad (8)$$

Acknowledgments. It has been a great pleasure and honour to collaborate with M.D. Scadron over many years. With this manuscript we would like to deliver our very best wishes to Mike and Arlene on the occasion of Mike's 70th birthday on February 12, 2008. This work has been supported by the FCT of the *Ministério da Ciência, Tecnologia e Ensino Superior* of Portugal, under Grants no. PDCT/FP/63907/2005, POCI/FP/81913/2007 and the Doppler and Nuclear Physics Institute (Dep. Theor. Phys.) at the Academy of Sciences of the Czech Republic by Project no. LC06002.

REFERENCES

1. F. Kleefeld, arXiv:0802.1540 [hep-ph].
2. F. Kleefeld, *Czech. J. Phys.* **55** (2005) 1123 [hep-th/0506140].
3. T. Eguchi, *Phys. Rev. D* **14** (1976) 2755.
4. M. Chaichian, R. Gonzalez Felipe, K. Huitu, *Phys. Lett. B* **363** (1995) 101 [hep-ph/9509223]; G. Kleppe, hep-th/9209124; N. G. Deshpande *et al.*, *Phys. Lett. B* **130** (1983) 61; *Phys. Rev. D* **29** (1984) 2851; T. Inami, H. Nishino, S. Watamura, *Phys. Lett. B* **117** (1982) 197; L. Castellani, P. Van Nieuwenhuizen, *Nucl. Phys. B* **213** (1983) 305; J. Kubo *et al.*, *Phys. Lett. B* **220** (1989) 191.
5. J.C. Collins, “*Renormalization*.”, ©1984 by Cambridge University Press.
6. R. Delbourgo, M. D. Scadron, *Mod. Phys. Lett. A* **10** (1995) 251 [hep-ph/9910242].
7. R. Delbourgo, M. D. Scadron, *Int. J. Mod. Phys. A* **13** (1998) 657 [hep-ph/9807504]; M. D. Scadron, hep-ph/0007184; M. D. Scadron, *Phys. Rev. D* **57** (1998) 5307 [hep-ph/9712425].
8. T. Hakioglu, M.D. Scadron, *Phys. Rev. D* **42** (1990) 941.
9. F. Kleefeld, *J. Phys. A* **39** (2006) L9 [hep-th/0506142]; X. Chen *et al.*, *Nucl. Phys. A* **759** (2005) 188.
10. F. Kleefeld, hep-th/0408028, hep-th/0408097.
11. F. Kleefeld, *AIP Conf. Proc.* **660** (2003) 325 [hep-ph/0211460].
12. F. Kleefeld, *AIP Conf. Proc.* **814** (2006) 690 [hep-ph/0510226].
13. C.M. Bender, S. Boettcher, *Phys. Rev. Lett.* **80** (1998) 5243; M. Znojil, *Rend. Circ. Mat. Palermo, Serie II, Suppl.* **72** (2004) 211 [math-ph/0104012]; C.M. Bender, *Rept. Prog. Phys.* **70** (2007) 947 [hep-th/0703096].
14. E. van Beveren *et al.*, *Mod. Phys. Lett. A* **17** (2002) 1673 [hep-ph/0204139]; M. D. Scadron, F. Kleefeld, G. Rupp, E. van Beveren, *Nucl. Phys. A* **724** (2003) 391 [hep-ph/0211275].

For convenience we have used the following short-hand notation for various relevant integrals [1]:

$$I_n(m^2) \equiv \int \frac{d^4 p}{(2\pi)^4} \frac{1}{(p^2 - m^2)^n} \stackrel{n \geq 3}{=} (-1)^n \frac{i}{16\pi^2} \frac{1}{(n-1)! m^{2n-4}}, \quad (5)$$

$$I_{n_1, n_2}(m_1^2, m_2^2) \equiv \int \frac{d^4 p}{(2\pi)^4} \frac{1}{(p^2 - m_1^2)^{n_1} (p^2 - m_2^2)^{n_2}}, \quad (6)$$

$$I_{n_1, n_2, n_3}^{sunset}(m_1^2, m_2^2, m_3^2) \equiv \int \frac{d^4 p_1}{(2\pi)^4} \int \frac{d^4 p_2}{(2\pi)^4} \int \frac{d^4 p_3}{(2\pi)^4} \frac{(2\pi)^4 \delta^4(p_1 + p_2 + p_3)}{(p_1^2 - m_1^2)^{n_1} (p_2^2 - m_2^2)^{n_2} (p_3^2 - m_3^2)^{n_3}}. \quad (7)$$

In all integrals we assume the imaginary part of the squared masses to be negative.




FIGURE 1. Tadpole sum: contributions to the σ one-point function




FIGURE 2. Quark mass: contributions to the quark self-energy




FIGURE 3. Sigma mass: contributions to the σ self-energy




FIGURE 4. Pion mass: contributions to the π self-energy